

Invariants of Legendrian Knots

Yu. V. Chekanov*

Abstract

We present two different constructions of invariants for Legendrian knots in the standard contact space $\mathbb{R}^3$. These invariants are defined combinatorially, in terms of certain planar projections, and are useful in distinguishing Legendrian knots that have the same classical invariants but are not Legendrian isotopic.

2000 Mathematics Subject Classification: 57R17.

Keywords and Phrases: Legendrian submanifold, Legendrian knot.

1. Introduction

1.1. Legendrian knots

A smooth knot L in the standard contact space $(\mathbb{R}^3, \alpha) = (\{(q, p, u)\}, du - pdq)$ is called Legendrian if it is everywhere tangent to the 2-plane distribution $\ker(\alpha)$ (or, in other words, if the restriction of α to L vanishes). Two Legendrian knots are Legendrian isotopic if they can be connected by a smooth path in the space of Legendrian knots (or, equivalently, if one can be sent to another by a diffeomorphism g of $\mathbb{R}^3$ such that $g^*\alpha = \varphi\alpha$, where $\varphi > 0$). In order to visualize a knot in $\mathbb{R}^3$, it is convenient to project it to a plane. In the Legendrian case, the character of the resulting picture will depend on the choice of the projection. The useful two are: the Lagrangian projection $\pi: \mathbb{R}^3 \rightarrow \mathbb{R}^2$, $(q, p, u) \mapsto (q, p)$, and the front projection $\sigma: \mathbb{R}^3 \rightarrow \mathbb{R}^2$, $(p, q, u) \mapsto (q, u)$. In Figure 1, two projections of the simplest Legendrian knot (unknot) are shown.

We say that a Legendrian knot $L \subset \mathbb{R}^3$ is π -generic if all self-intersections of the immersed curve $\pi(L)$ are transverse double points. We can represent a π -generic Legendrian knot L by its (Lagrangian) diagram: the curve $\pi(L) \subset \mathbb{R}^2$, at every crossing of which the overpassing branch (the one with the greater value of u) is marked. Of course, not every abstract knot diagram in $\mathbb{R}^2$ is a diagram of

*Moscow Center for Continuous Mathematical Education, B. Vlasievsky per. 11, Moscow 119002, Russia. E-mail: chekanov@mccme.ru




Figure 1: Lagrangian projection and front projection

a Legendrian knot, or is oriented diffeomorphic to such (it requires a bit of extra work to check whether a given diagram corresponds to a Legendrian knot, cf. [1]).

Given a Legendrian knot $L \subset \mathbb{R}^3$, its σ -projection, or front, $\sigma(L) \subset \mathbb{R}^2$ is a singular curve with nowhere vertical tangent vectors. Its singularities, generically, are semi-cubic cusps and transverse double points. We say that L is σ -generic if, moreover, all self-intersections of $\sigma(L)$ have different q -coordinates. Every closed planar curve with these types of singularities and nowhere vertical tangent vectors is a front of a Legendrian knot. Note that there is no need to explicitly indicate the type of a crossing of a front: the overpassing branch (the one with the greater value of p) is always the one with the greater slope.

1.2. Classical invariants

The so-called classical invariants of an oriented Legendrian knot L are defined as follows. The first of them is, formally, just the smooth isotopy type of L . The Thurston–Bennequin number $\beta(L)$ of L is the linking number (with respect to the orientation defined by $\alpha \wedge d\alpha$) between L and $s(L)$, where s is a small shift along the u direction. The Maslov number $m(L)$ (which actually is an invariant of Legendrian immersion) is twice the rotation number of the projection of L to the (q, p) plane (or, equivalently, the value of the Maslov 1-cohomology class on the fundamental class of L). The change of orientation on L changes the sign of $m(L)$ and preserves $\beta(L)$. The Thurston–Bennequin number of a π -generic Legendrian knot L can be computed by counting the crossings of its Lagrangian diagram $\pi(L)$ with signs:

$$\beta(L) = \#(\nearrow \searrow) - \#(\nwarrow \swarrow)$$

(where the q axis is horizontal and the p axis is vertical). In terms of the front projection, the classical invariants can be computed as follows. The Maslov number of a σ -generic oriented Legendrian knot L is the number of the right cusps of the front $\sigma(L)$, counted with signs depending on the orientations:

$$m(L) = \#(\nearrow) - \#(\searrow).$$

The Thurston–Bennequin number of L is the number of crossings of $\sigma(L)$ counted with signs minus half the total number of cusps (= the number of right cusps):

$$\beta(L) = \#(\nearrow \searrow) + \#(\nwarrow \swarrow) - \#(\nearrow \nwarrow) - \#(\searrow \swarrow) - \#(\nearrow) - \#(\searrow).$$

For the Legendrian knot shown in Figure 1, we have $m = 0$, $\beta = -1$.

1.3. New invariants and classification results

It is easy to show that every smooth knot admits a Legendrian realization. A natural question to ask is whether there exists a pair of Legendrian knots which have the same classical invariants but are not Legendrian isotopic. The answer is positive:

Theorem 1.1. [1, 2] *There exist Legendrian knots L, L' (see Figure 2 on p. 390, Figure 6 on p. 393) that have the same classical invariants (smooth knot type 5_2 , $m = 0$, $\beta = 1$) but are not Legendrian isotopic.*

In the next two sections, we present two combinatorial constructions of Legendrian knot invariants. The first one associates to the Lagrangian projection of a Legendrian knot a differential graded algebra (DGA). The second construction deals with decompositions of the front projection into closed curves. Each of the two provides a proof of Theorem 1.1. The invariants do not change when the orientation of the knot reverses, so essentially they are invariants of non-oriented Legendrian knots. It should be mentioned that these constructions also produce, with minor modifications, invariants of Legendrian links.

The number of Legendrian knots with given classical invariants is known to be finite [3]. Eliashberg and Fraser gave a classification of Legendrian realization for smooth unknots [5, 6]. It turned out that smooth unknots are Legendrian simple in the sense that the Legendrian isotopy types of their Legendrian realizations are determined by the classical invariants. Etnyre and Honda obtained a classification of Legendrian realization for torus knots and the figure eight knot [8]. Again, these smooth knot types proved to be Legendrian simple. The 5_2 is the simplest knot type for which the classification is not known. By Theorem 1.1, the type 5_2 is not Legendrian simple. Conjecturally, two Legendrian knots of smooth type 5_2 with the same classical invariants are Legendrian isotopic unless they form the pair L, L' from Theorem 1.1. Several interesting examples of knots with coinciding classical invariants but not Legendrian isotopic were constructed by Ng [14, 15].

2. DGA of a Legendrian knot

2.1. Definitions

In this section, we associate with every π -generic Legendrian knot L a DGA (A, ∂) over $\mathbb{Z}/2\mathbb{Z}$ ([1]; a similar construction was also given by Eliashberg). This DGA is related to the symplectic field theory introduced by Eliashberg, Givental, and Hofer in [7] (see [10]).

Let $\{a_1, \dots, a_n\}$ be the set of crossings of $Y = \pi(L)$. Define A to be the tensor algebra (free associative unital algebra) $T(a_1, \dots, a_n)$ with generators $a_1, \dots, a_n$. The grading on A takes values in the group $\mathbb{Z}/m(L)\mathbb{Z}$ and is defined as follows. Given a crossing a_j , consider the points $z_+, z_- \in L$ such that $\pi(z_+) = \pi(z_-) = a_j$ and the u -coordinate of z_+ is greater than the u -coordinate of z_- . These points divide L into two pieces, γ_1 and γ_2 , which we orient from z_+ to z_- . We can assume,

without loss of generality, that the intersecting branches are orthogonal at a . Then, for $\varepsilon \in \{1, 2\}$, the rotation number of the curve $\pi(\gamma_\varepsilon)$ has the form $N_\varepsilon/2 + 1/4$, where $N_\varepsilon \in \mathbb{Z}$. Clearly, $N_1 - N_2$ is equal to $\pm m(L)$. Hence N_1 and N_2 represent the same element of the group $\Gamma = \mathbb{Z}/m(L)\mathbb{Z}$, which we define to be the degree of a_j .

We are going to define the differential ∂ . For every natural k , fix a (curved) convex k -gon $\Pi_k \subset \mathbb{R}^2$ whose vertices $x_0^k, \dots, x_{k-1}^k$ are numbered counter-clockwise. The form $dq \wedge dp$ defines an orientation on $\mathbb{R}^2$. Denote by $W_k(Y)$ the collection of smooth orientation-preserving immersions $f: \Pi_k \rightarrow \mathbb{R}^2$ such that $f(\partial\Pi_k) \subset Y$. Note that $f \in W_k(Y)$ implies $f(x_i^k) \in \{a_1, \dots, a_n\}$. Consider the set of nonparametrized immersions $\widetilde{W}_k(Y)$, which is the quotient of $W_k(Y)$ by the action of the group $\{g \in \text{Diff}_+(\Pi_k) \mid g(x_i^k) = x_i^k\}$. The diagram Y divides a neighbourhood of each of its crossings into four sectors. We call positive two of them which are swept out by the underpassing curve rotating counter-clockwise, and negative the other two (the sectors are marked in Figure 1). For each vertex x_i^k of the polygon Π_k , a smooth immersion $f \in \widetilde{W}_k(Y)$ maps its neighbourhood in Π_k to either a positive or a negative sector; we shall say that x_i^k is, respectively, a positive or a negative vertex for f . Define the set $W_k^+(Y)$ to consist of immersions $f \in \widetilde{W}_k(Y)$ such that the vertex x_0^k is positive for f , and all other vertices are negative. Let $W_k^+(Y, a_j) = \{f \in W_k^+(Y) \mid f(x_0^k) = a_j\}$. Denote $A_1 = \{a_1, \dots, a_n\} \otimes \mathbb{Z}/2\mathbb{Z} \subset A$, $A_k = (A_1)^{\otimes k}$. Then $A = \oplus_{l=0}^\infty A_l$. Let $\partial = \sum_{k \geq 0} \partial_k$, where $\partial_k(A_i) \in A_{i+k-1}$. Define

$$\partial_k(a_j) = \sum_{f \in W_{k+1}^+(Y, a_j)} f(x_1) \cdots f(x_k)$$

(for $k = 0$, we have $\partial_0(a_j) = \#(W_1^+(Y, a_j))$, and extend ∂ to A by linearity and the Leibniz rule. The following theorem says that (A, ∂) is indeed a DGA:

Theorem 2.1. *The differential ∂ is well defined. We have $\deg(\partial) = -1$ and $\partial^2 = 0$.*

Define the (l -th, where $l \in \Gamma$) stabilization of a DGA $(T(a_1, \dots, a_n), \partial)$ to be the DGA $(T(a_1, \dots, a_n, a_{n+1}, a_{n+2}), \partial)$, where $\deg(a_{n+1}) = l$, $\deg(a_{n+2}) = l - 1$, $\partial(a_{n+1}) = a_{n+2}$, and ∂ acts on $a_1, \dots, a_n$ as before. An automorphism of $T(a_1, \dots, a_n)$ is called elementary if it sends a_i to $a_i + v$, where v does not involve a_i , and fixes a_j for $j \neq i$. Two DGAs $(T(a_1, \dots, a_n), \partial)$, $(T(a_1, \dots, a_n), \partial')$ are called tame isomorphic if one can be obtained from another by a composition of elementary automorphisms; they are called stable tame isomorphic if they become tame isomorphic after (iterated) stabilizations.

Theorem 2.2. *Let (A, ∂) , (A', ∂') be the DGAs of $(\pi$ -generic) Legendrian knots L, L' . If L and L' are Legendrian isotopic then (A, ∂) and (A', ∂') are stable tame isomorphic. In particular, the homology rings $H(A, \partial) = \ker(\partial)/\text{im}(\partial)$ and $H(A', \partial') = \ker(\partial')/\text{im}(\partial')$ are isomorphic as graded rings.*

The hard part in the proof of Theorem 2.1 is to show that $\partial^2 = 0$. The proof of this fact mimics, in a combinatorial way, the classical gluing–compactness argument of the Floer theory (cf. [11]). The proof of Theorem 2.2 involves a careful study of

the behaviour of the DGA associated with a Legendrian knot when its Lagrangian diagram goes through elementary bifurcations (Legendrian Reidemeister moves).

It turns out that one cannot replace the coefficient ring $\mathbb{Z}/2\mathbb{Z}$ by $\mathbb{Z}$: in some sense, our homology theory is not oriented. However, the construction described above can be modified to associate with a Legendrian knot L a DGA graded by $\mathbb{Z}$ and having $\mathbb{Z}[s, s^{-1}]$ (where $\deg(s) = m(L)$) as a coefficient ring [10]. After reducing the grading to $\mathbb{Z}/m(L)\mathbb{Z}$, and applying the homomorphism $\mathbb{Z}[s, s^{-1}] \rightarrow \mathbb{Z}/2\mathbb{Z}$ sending both s and $1 \in \mathbb{Z}$ to $1 \in \mathbb{Z}/2\mathbb{Z}$, this $\mathbb{Z}[s, s^{-1}]$ -DGA becomes the $\mathbb{Z}/2\mathbb{Z}$ -DGA of the knot L .

2.2. Poincaré polynomials

Homology rings of DGAs can be hard to work with. We are going to define an easily computable invariant I , which is a finite subset of the group monoid $\mathbb{N}_0[\Gamma]$, where $\mathbb{N}_0 = \{0, 1, \dots\}$, $\Gamma = \mathbb{Z}/m(L)\mathbb{Z}$. Assume that $\partial_0 = 0$. Then $\partial_1^2 = 0$. Since $\partial(A_1) \subset A_1$, we can consider the homology $H(A_1, \partial_1) = \ker(\partial_1|_{A_1})/\text{im}(\partial_1|_{A_1})$, which is a vector space graded by the cyclic group Γ . Define the Poincaré polynomial $P_{(A, \partial)} \in \mathbb{N}_0[\Gamma]$ by

$$P_{(A, \partial)}(t) = \sum_{\lambda \in \Gamma} \dim(H_\lambda(A_1, \partial_1)) t^\lambda,$$

where $H_\lambda(A_1, \partial_1)$ is the degree λ homogeneous component of $H(A_1, \partial_1)$. Define the group $\text{Aut}_0(A)$ to consist of graded automorphisms of A such that for each $i \in \{1, \dots, n\}$ we have $g(a_i) = a_i + c_i$, where $c_i \in A_0 = \mathbb{Z}/2\mathbb{Z}$. (of course, $c_i = 0$ when $\deg(a_i) \neq 0$). Consider the set $U_0(A, \partial)$ consisting of automorphisms $g \in \text{Aut}_0(A)$ such that $(\partial^g)_0 = 0$ (where $\partial^g = g^{-1} \circ \partial \circ g$). Define

$$I(A, \partial) = \{P_{(A, \partial^g)} \mid g \in U_0(A, \partial)\}.$$


Since $\text{Aut}_0(A)$ has at most 2^n elements, this invariant is not hard to compute. We can associate with every (π -generic) Legendrian knot L the set $I(L) = I(A_L, \partial_L)$. Note that $P(-1) = \beta(L)$ for $P \in I(L)$. One can show that I is an invariant of stable tame DGA isomorphism. Hence Theorem 2.2 implies the following

Corollary 2.3. *If L is Legendrian isotopic to L' then $I(L) = I(L')$.*

The set $I(L)$ can be empty (cf. Section 4) but no examples are known where $I(L)$ contains more than one element. Also, for all known examples of pairs L, L' of Legendrian knots with coinciding classical invariants we have $P(1) = P'(1)$, where $P \in I(L)$, $P' \in I(L')$. Other, more complicated invariants of stable tame isomorphism were developed and applied to distinguishing Legendrian knots in [15].


2.3. Examples

1a. Let $(A, \partial) = (T(a_1, \dots, a_9), \partial)$ be the DGA of the Legendrian knot L given in Figure 2. We have $m(L) = 0$, $\beta(L) = 1$, $\deg(a_i) = 1$ for $i \leq 4$, $\deg(a_5) = 2$, $\deg(a_6) = -2$, $\deg(a_i) = 0$ for $i \geq 7$, $\partial(a_1) = 1 + a_7 + a_7 a_6 a_5$, $\partial(a_2) = 1 + a_9 + a_5 a_6 a_9$, $\partial(a_3) = 1 + a_8 a_7$, $\partial(a_4) = 1 + a_8 a_9$, $\partial(a_i) = 0$ for $i \geq 5$.

Figure 2: Lagrangian projections of two Legendrian 5_2 knots

1b. Let $(A', \partial) = (T(a_1, \dots, a_9), \partial)$ be the DGA of the Legendrian knot L' given in Figure 2. We have $m(L') = 0$, $\beta(L') = 1$, $\deg(a_i) = 1$ for $i \leq 4$, $\deg(a_i) = 0$ for $i \geq 5$, $\partial(a_1) = 1 + a_7 + a_5 + a_7a_6a_5 + a_9a_8a_5$, $\partial(a_2) = 1 + a_9 + a_5a_6a_9$, $\partial(a_3) = 1 + a_8a_7$, $\partial(a_4) = 1 + a_8a_9$, $\partial(a_i) = 0$ for $i \geq 5$.

An explicit computation shows that $I(L) = \{t^{-2} + t^1 + t^2\}$, $I(L') = \{2t^0 + t^1\}$ and hence Theorem 1.1 follows from Corollary 2.3.

Figure 3: Lagrangian projection of a Legendrian 6_2 knot

2. [14] Let $(A, \partial) = (T(a_1, \dots, a_{11}), \partial)$ be the DGA of the Legendrian knot K given in Figure 3. We have $m(K) = 0$, $\beta(K) = -7$, $\deg(a_i) = 1$ for $i \in \{1, 2, 7, 9, 10\}$, $\deg(a_i) = 0$ for $i \in \{3, 4\}$, $\deg(a_i) = -1$ for $i \in \{5, 6, 8, 11\}$; $\partial(a_1) = 1 + a_{10}a_5a_3$, $\partial(a_2) = 1 + a_3 + a_3a_6a_{10} + a_3a_{11}a_7$, $\partial(a_4) = a_5 + a_{11} + a_{11}a_7a_5$, $\partial(a_6) = a_{11}a_8$, $\partial(a_7) = a_8a_{10}$, $\partial(a_9) = 1 + a_{10}a_{11}$, $\partial(a_i) = 0$ for $i \in \{3, 5, 8, 10, 11\}$. Denote by $\widehat{K}$ the ‘Legendrian mirror’ of K — the image of K under the map $(q, p, u) \mapsto (-q, p, -u)$. The Legendrian knots $K, \widehat{K}$ have the same classical invariants. However, they are not Legendrian isotopic, and it is possible to distinguish them by means of their DGAs. There exist homology classes ξ_+, ξ_- in the graded homology ring $H(A, \partial)$ such that $\deg(\xi_+) = 1$, $\deg(\xi_-) = -1$, and $\xi_+ \xi_- = 1$ (choose $\xi_+ = [a_{10}]$, $\xi_- = [a_{11}]$). It follows from the definitions that the DGA for $\widehat{K}$ is obtained from (A, ∂) by applying the anti-automorphism reversing the order of gener-

ators in all monomials. Thus, if K and $\widehat{K}$ are Legendrian isotopic then the graded homology ring $H(A, \partial)$ is anti-isomorphic to itself, and there exist $\xi'_+, \xi'_- \in H(A, \partial)$ such that $\deg(\xi'_+) = 1$, $\deg(\xi'_-) = -1$, $\xi'_- \xi'_+ = 1$. But one can check that such classes do not exist (see [14, 15] for details) and hence K and $\widehat{K}$ are not Legendrian isotopic. Note that ‘first order invariants’ such as Poincaré polynomials are useless in distinguishing Legendrian mirror knots.

3. Admissible decompositions of fronts

3.1. Definitions

In this section, we present the invariants of Legendrian knots constructed in [2]. These invariants are defined in terms of the front projection.

Given a σ -generic oriented Legendrian knot L , denote by $C(L)$ the set of its points corresponding to cusps of $\sigma(L)$. The Maslov index $\mu: L \setminus C(L) \rightarrow \Gamma = \mathbb{Z}/m(L)\mathbb{Z}$ is a locally constant function, uniquely defined up to an additive constant by the following rule: the value of μ jumps at points of $C(L)$ by ± 1 as shown in Figure 4. We call a crossing of $\Sigma = \sigma(L)$ Maslov if μ takes the same value on both its branches.




Figure 4: Jumps of the Maslov index near cusps

Assume that $\Sigma = \sigma(L)$ is a union of closed curves $X_1, \dots, X_n$ that have finitely many self-intersections and meet each other at finitely many points. Then we call the unordered collection $\{X_1, \dots, X_n\}$ a decomposition of Σ . A decomposition $\{X_1, \dots, X_n\}$ is called admissible if it satisfies certain conditions, which we are going to define. The first two are as follows:

- (1) Each curve X_i bounds a topologically embedded disk: $X_i = \partial B_i$.
- (2) For each $i \in \{1, \dots, n\}$, $q \in \mathbb{R}$, the set $B_i(q) = \{u \in \mathbb{R} \mid (q, u) \in B_i\}$ is either a segment, or consists of a single point u such that (q, u) is a cusp of Σ , or is empty.

Conditions (1) and (2) imply that each curve X_i has exactly two cusps (and hence the number of curves is half the number of cusps). Each X_i is divided by cusps into two pieces, on which the coordinate q is a monotone function. Near a crossing $x \in X_i \cap X_j$, the decomposition of Σ may look in one of the three ways represented in Figure 5. Conditions (1) and (2), in particular, rule out the decomposition shown in Figure 5a. We call the crossing point x switching if X_i and X_j are not smooth near x (Figure 5b), and non-switching otherwise (Figure 5c).

- (3) If $(q_0, u) \in X_i \cap X_j$ ($i \neq j$) is switching then for each $q \neq q_0$ sufficiently close to q_0 the set $B_i(q) \cap B_j(q)$ either coincides with $B_i(q)$ or $B_j(q)$, or is empty.




Figure 5: Local decompositions

- (4) Every switching crossing is Maslov.

We call a decomposition admissible if it satisfies Conditions (1)-(3), and graded admissible if it also satisfies Condition (4). Denote by $\text{Adm}(\Sigma)$ (resp. $\text{Adm}_+(\Sigma)$) the set of admissible (resp. graded admissible) decompositions of Σ . Given $D \in \text{Adm}(\Sigma)$, denote by $\text{Sw}(D)$ the set of its switching points. Define $\theta(D) = \#(D) - \#(\text{Sw}(D))$.

Theorem 3.1. *If σ -generic Legendrian knots $L, L' \subset \mathbb{R}^3$ are Legendrian isotopic then there exists a one-to-one mapping $g: \text{Adm}(\sigma(L)) \rightarrow \text{Adm}(\sigma(L'))$ such that $g(\text{Adm}_+(\sigma(L))) = \text{Adm}_+(\sigma(L'))$ and $\theta(g(D)) = \theta(D)$ for each $D \in \text{Adm}(\sigma(L))$. In particular, the numbers $\#(\text{Adm}(\sigma(L)))$ and $\#(\text{Adm}_+(\sigma(L)))$ are invariants of Legendrian isotopy.*


3.2. Remarks

1. Decompositions of fronts were first considered by Eliashberg in [4] (only Conditions (1) and (2) were involved).
2. No examples are known where the total number of admissible decompositions $\#(\text{Adm}(\Sigma))$ is different for two Legendrian knots with coinciding classical invariants.
3. The proof of Theorem 3.1 goes as follows: we connect L with L' by a generic path in the space of Legendrian knots and define a canonical way to extend admissible decompositions through the points where the front is not σ -generic. The mapping g depends on the choice of the path: a loop in the space of Legendrian knots lifts to an automorphism of $\text{Adm}(\sigma(L))$ which can be non-trivial even when the loop is contractible. The meaning of this phenomenon is not clear.
4. It would be interesting to understand the relation between admissible decompositions and DGAs of Legendrian knots. The first result in this direction is that if $\text{Adm}_+(\sigma(L))$ is nonempty then the set $I(L)$ defined in the previous section is also nonempty [12].

3.3. Examples

Note that every admissible decomposition D of a front Σ is uniquely defined by its set of switching points. Indeed, denote by $X(\Sigma)$ the set of crossings of Σ , then each subset $E \subset X(\Sigma)$ defines a decomposition $D(E)$ of Σ which near $x \in X(\Sigma)$ has the form shown in Figure 5b if $x \in E$, and the form shown in Figure 5c otherwise. Clearly, if $E = \text{Sw}(D)$ then $D = D(E)$.

The Legendrian knots represented by the fronts Σ, Σ' in Figure 6 are respectively Legendrian isotopic to the Legendrian knots L, L' defined in Figure 2. We are

Figure 6: Fronts of two Legendrian 5_2 knots

going to show that $\#(\text{Adm}_+(\Sigma)) = 1$, $\#(\text{Adm}_+(\Sigma')) = 2$, and hence Theorem 1.1 is a consequence of Theorem 3.1. Assume that $D \in \text{Adm}(\Sigma)$. Consider the curve $X_1 \in D$ containing the piece of Σ indicated by the lower arrow. Being applied to X_1 , Conditions (1) and (2) imply that $c_2, c_3 \in \text{Sw}(D)$. Similarly, looking at the curve $X_2 \in D$ containing the piece of Σ indicated by the upper arrow, we conclude that $c_4, c_5 \in \text{Sw}(D)$. If one of the crossings c_1, c_6 is switching, so is the other. Then either $\text{Sw}(D) = \{c_2, c_3, c_4, c_5\}$ or $\text{Sw}(D) = \{c_1, c_2, c_3, c_4, c_5, c_6\}$. It is not hard to check that both decompositions are admissible but only the first one is graded. Thus $\#(\text{Adm}_+(\Sigma)) = 1$. Arguing similarly, one can find that $\#(\text{Adm}(\Sigma')) = 2$, where the admissible decompositions D_1, D_2 are defined by $\text{Sw}(D_1) = \{c_2, c_3, c_4, c_5\}$, $\text{Sw}(D_2) = \{c_1, c_2, c_3, c_4, c_5, c_6\}$, and are both graded.

4. Instability of invariants

There are two stabilizing operations, S_- and S_+ , on Legendrian isotopy classes of oriented Legendrian knots, defined as follows. Given an oriented Legendrian knot L , we perform one of the operations shown in Figure 7 in a small neighbourhood of a point on L . One can check that, up to Legendrian isotopy, the resulting Legendrian knot $S_{\pm}(L)$ does not depend on the choices involved, and the operations S_-, S_+ commute. An important observation is that two Legendrian knots L, L' have the same classical invariants if and only if they are stable Legendrian isotopic in the sense that there exist $n_-, n_+ \in \mathbb{N}_0$ such that $S_-^{n_-}(S_+^{n_+}(L))$ is Legendrian isotopic to $S_-^{n_-}(S_+^{n_+}(L'))$ [13].




Figure 7: Stabilizations

Thus the invariants constructed in Sections 2 and 3 cannot be stable. In fact, they fail already after the first stabilization. The homology ring H of the DGA corresponding to $S_{\pm}(L)$ vanish, and the set $I(S_{\pm}(L))$ is empty. This can be easily

derived from the fact that the DGA of $S_{\pm}(L)$ can be obtained from the DGA of L by adding a new generator a such that $\partial(a) = 1$. The front of $S_{\pm}(L)$ has no admissible decompositions because Conditions (1) and (2) cannot hold for the curve X_i containing the newly created cusps.

Studying Legendrian realizations of non-prime knots, Etnyre and Honda constructed, for each m , examples of Legendrian knots that have the same classical invariants but are not Legendrian isotopic even after m stabilizations [9]. Their proof uses the classification of Legendrian torus knots given in [8]. It is an open problem to find invariants distinguishing those knots, or any pair of stabilized knots with the same classical invariants.

References

- [1] Yu. V. Chekanov, Differential algebra of Legendrian links, to appear in *Inventiones Mathematicae*.
- [2] Yu. V. Chekanov & P. E. Pushkar, in preparation.
- [3] V. Colin, E. Giroux & K. Honda, On the coarse classification of tight contact structures, Preprint, 2002.
- [4] Ya. Eliashberg, A theorem on the structure of wave fronts and its application in symplectic topology, *Funct. Anal. Appl.*, 21 (1987), 227–232.
- [5] Ya. Eliashberg, Legendrian and transversal knots in tight contact 3-manifolds, In: *Topological methods in modern mathematics (Stony Brook, NY, 1991)*, Publish or Perish, 1993, 171–193.
- [6] Ya. Eliashberg & M. Fraser, Classification of topologically trivial Legendrian knots, In: *Geometry, topology, and dynamics (Montreal, PQ, 1995)*, CRM Proc. Lecture Notes, 15, AMS, Providence, 1998, 17–51.
- [7] Ya. Eliashberg, A. Givental, & H. Hofer, An introduction to symplectic field theory, *Geom. Funct. Anal.* (2000), Special Volume, Part II, 560–673.
- [8] J. Etnyre & K. Honda, Knots and contact geometry I: torus knots and the figure eight knot, Preprint, 2000, math.GT/0006112.
- [9] J. Etnyre & K. Honda, Knots and Contact Geometry II: Connected Sums, Preprint, 2002, math.GT/0205310.
- [10] J. Etnyre, L. Ng, & J. Sabloff, Invariants of Legendrian knots and coherent orientations, Preprint, 2001, math.GT/0101145.
- [11] A. Floer, Symplectic fixed points and holomorphic spheres, *Comm. Math. Phys.*, 120, 1989, 575–611.
- [12] D. Fuchs, Private communication, 2001.
- [13] D. Fuchs & S. Tabachnikov, Invariants of Legendrian and transverse knots in the standard contact space, *Topology*, 36 (1997), 1025–1053.
- [14] L. Ng, Legendrian mirrors and Legendrian isotopy, Preprint, 2000, math.GT/0008210.
- [15] L. Ng, Computable Legendrian invariants, Preprint, 2001, math.GT/0011265.